

Seizure Rescue Medications



NAYZILAM

Midazolam Nasal Spray

Approved for ages 12 years and older



DO NOT PRESS THE PLUNGER YET

Hold the nasal spray device with your thumb on the plunger and your middle and index fingers on each side of the nozzle.

If you press the plunger now, you will lose the medicine.



DO NOT PRESS THE PLUNGER YET

Place the tip of the nozzle into one nostril until your fingers are against the bottom of the patient's nose.



PRESS

Press the plunger firmly.

How to use
Nayzilam.



VALTOCO

Diazepam Nasal Spray

Approved for ages 6 years and older



DO NOT PRESS THE PLUNGER YET

Hold the nasal spray device with your thumb. If your child needs either 15mg or 20mg of Valtoco, use both devices, one in each nostril to give them the right amount.



DO NOT PRESS THE PLUNGER YET

Place the tip of the nozzle into one nostril until your fingers are against the bottom of the patient's nose.



PRESS

Press the plunger firmly.

How to use
Valtoco.



DIASTAT

Diazepam Nasal Gel

Approved for ages 2 years and older

Diastat comes as a gel inside prefilled syringes that are used to administer the medication rectally.

1. Put person on their side where they can't fall.
2. Get medicine. Get syringe. Note: Seal pin is attached to the cap.
3. Push up with thumb and pull to remove cap from syringe. Be sure seal pin is removed with the cap.
4. Lubricate rectal tip with lubricating jelly.
5. Turn person on side facing you.
6. Bend upper leg forward to expose rectum.
7. Separate buttocks to expose rectum.
8. Gently insert syringe tip into rectum. Note: Rim should be snug against rectal opening.
9. Slowly count to 3 while gently pushing plunger in until it stops.
10. Slowly count to 3 before removing syringe from rectum.
11. Slowly count to 3 while holding buttocks together to prevent leakage.
12. Keep person on the side facing you, note time given and continue to observe.

How to use
Diastat.



Savings and Assistance

You may qualify for savings and assistance for these medications. For more information, please scan the QR code below.

Nayzilam[®]
(midazolam) nasal spray

For Savings



VALTOCO[®]
(diazepam nasal spray) ©



Seizure First Aid

IF YOUR CHILD HAS A SEIZURE:

- Stay calm and track the time.
- Turn them on their left side.
- Remove nearby objects to prevent injury.
- DON'T try to stop the shaking.
- DON'T put anything into your child's mouth.
- Stay with your child until they are fully conscious.



Watch Our Training Video

Seizure Emergency

CALL 911 IF:

- Your child has a seizure for the first time.
- The seizure lasts longer than five minutes, or seizures keep happening.
- Your child stays unconscious after a seizure.
- The seizure happens in water.
- Your child gets injured or has trouble breathing.
- Your child is diabetic.